

Aequat. Math. 88 (2014), 163–168

© The Author(s) 2013. This article is published

with open access at Springerlink.com

0001-9054/14/010163-6

published online June 15, 2013

DOI 10.1007/s00010-013-0214-x

## Remark on hyperstability of the general linear equation

MAGDALENA PISZCZEK

**Abstract.** We present a result concerning the hyperstability of the general linear equation. Namely, we show that a function satisfying the equation approximately must be actually a solution to it.

**Mathematics Subject Classification (2010).** Primary 39B82, 39B62; Secondary 47H14, 47H10.

**Keywords.** Hyperstability, general linear equation.

Throughout the paper  $\mathbb{F}$ ,  $\mathbb{K}$  denote the fields of real or complex numbers. Let  $X$ ,  $Y$  be normed spaces over  $\mathbb{F}$ ,  $\mathbb{K}$ , respectively. The aim of the paper is to give a hyperstability result for the general linear equation

$$g(ax + by) = Ag(x) + Bg(y), \quad (1)$$

where  $g: X \rightarrow Y$  and  $a, b \in \mathbb{F} \setminus \{0\}$ ,  $A, B \in \mathbb{K}$ . We see that for  $a = b = A = B = 1$  in (1) we get the Cauchy equation while the Jensen equation corresponds to  $a = b = A = B = \frac{1}{2}$ . The general linear equation has been studied by many authors, see for example [6, 8, 10–12, 14].

Probably the first hyperstability result was published in [3] and concerned ring homomorphisms. However the term hyperstability was used for the first time in [13].

We recall the most classical results concerning the Hyers-Ulam stability of the Cauchy equation

$$f(x + y) = f(x) + f(y), \quad x, y \in X, \quad (2)$$

which were the motivation to write the paper.

If  $X$  is a normed space,  $Y$  is a Banach space,  $c \geq 0$ ,  $p$  is a real number such that  $0 < p < 1$  and  $f: X \rightarrow Y$  satisfies the inequality

$$\|f(x + y) - f(x) - f(y)\| \leq c(\|x\|^p + \|y\|^p), \quad x, y \in X, \quad (3)$$

then there exists a unique solution  $T: X \rightarrow Y$  of (2) with

$$\|f(x) - T(x)\| \leq \frac{c\|x\|^p}{|1 - 2^{p-1}|}, \quad x \in X.$$

It is due to Aoki [1] (see also [15]). Later Gajda [7] obtained this result for  $p > 1$  and Rassias [16] for  $p < 0$ . Moreover, in [7] we can find an example showing that an analogous result for  $p = 1$  is not true. For  $p = 0$  it is the first result of stability proved by Hyers [9]. Now, it is known that for  $p < 0$  we have the hyperstability result, that is  $f$  satisfying (3) must be additive for  $x, y \in X \setminus \{0\}$  (see [4]).

The method of the proof used in the main theorem is based on a fixed point result that can be derived from [5] (Theorem 1). To present it we need the following three hypothesis:

- (H1)  $U$  is a nonempty set,  $Y$  is a Banach space,  $f_1, \dots, f_k: U \rightarrow U$  and  $L_1, \dots, L_k: U \rightarrow \mathbb{R}_+$  are given.  
 (H2)  $\mathcal{T}: Y^U \rightarrow Y^U$  is an operator satisfying the inequality

$$\|\mathcal{T}\xi(x) - \mathcal{T}\mu(x)\| \leq \sum_{i=1}^k L_i(x) \|\xi(f_i(x)) - \mu(f_i(x))\|,$$

$$\xi, \mu \in Y^U, x \in U.$$

- (H3)  $\Lambda: \mathbb{R}_+^U \rightarrow \mathbb{R}_+^U$  is defined by

$$\Lambda\delta(x) := \sum_{i=1}^k L_i(x) \delta(f_i(x)), \quad \delta \in \mathbb{R}_+^U, x \in U.$$

Now we are in a position to present the above mentioned fixed point theorem.

**Theorem 1.** *Let hypotheses (H1)–(H3) be valid and functions  $\varepsilon: U \rightarrow \mathbb{R}_+$  and  $\varphi: U \rightarrow Y$  fulfil the following two conditions*

$$\|\mathcal{T}\varphi(x) - \varphi(x)\| \leq \varepsilon(x), \quad x \in U,$$

$$\varepsilon^*(x) := \sum_{n=0}^{\infty} \Lambda^n \varepsilon(x) < \infty, \quad x \in U.$$

Then there exists a unique fixed point  $\psi$  of  $\mathcal{T}$  with

$$\|\varphi(x) - \psi(x)\| \leq \varepsilon^*(x), \quad x \in U.$$

Moreover

$$\psi(x) := \lim_{n \rightarrow \infty} \mathcal{T}^n \varphi(x), \quad x \in U.$$

**Theorem 2.** *Let  $X$  be a normed space over a field  $\mathbb{F}$ ,  $Y$  be a Banach space over  $\mathbb{K}$ ,  $a, b \in \mathbb{F} \setminus \{0\}$ ,  $A, B \in \mathbb{K}$ ,  $c \geq 0$ ,  $p < 0$  and  $g: X \rightarrow Y$  satisfy*

$$\|g(ax + by) - Ag(x) - Bg(y)\| \leq c(\|x\|^p + \|y\|^p), \quad x, y \in X \setminus \{0\}. \quad (4)$$

Then  $g$  satisfies the equation

$$g(ax + by) = Ag(x) + Bg(y), \quad x, y \in X \setminus \{0\}. \quad (5)$$

*Proof.* Replacing  $x$  by  $\frac{1}{a}(m+1)x$  and  $y$  by  $-\frac{1}{b}mx$  for  $m \in \mathbb{N}$  in (4) we get

$$\begin{aligned} & \left\| g(x) - Ag\left(\frac{1}{a}(m+1)x\right) - Bg\left(-\frac{1}{b}mx\right) \right\| \\ & \leq c \left( \left| \frac{1}{a}(m+1) \right|^p + \left| \frac{1}{b}m \right|^p \right) \|x\|^p, \quad x \in X \setminus \{0\}. \end{aligned} \quad (6)$$

Further put

$$\begin{aligned} \mathcal{T}_m \xi(x) &:= A\xi\left(\frac{1}{a}(m+1)x\right) + B\xi\left(-\frac{1}{b}mx\right), \quad x \in X \setminus \{0\}, \xi \in Y^{X \setminus \{0\}}, \\ \varepsilon_m(x) &:= c \left( \left| \frac{1}{a}(m+1) \right|^p + \left| \frac{1}{b}m \right|^p \right) \|x\|^p, \quad x \in X \setminus \{0\}. \end{aligned}$$

Then the inequality (6) takes the form

$$\|\mathcal{T}_m g(x) - g(x)\| \leq \varepsilon_m(x), \quad x \in X \setminus \{0\}.$$

The operator

$$\Lambda_m \eta(x) := |A|\eta\left(\frac{1}{a}(m+1)x\right) + |B|\eta\left(-\frac{1}{b}mx\right), \quad \eta \in \mathbb{R}_+^{X \setminus \{0\}}, x \in X \setminus \{0\}$$

has the form described in (H3) with  $k = 2$  and  $f_1(x) = \frac{1}{a}(m+1)x$ ,  $f_2(x) = -\frac{1}{b}mx$ ,  $L_1(x) = |A|$ ,  $L_2(x) = |B|$  for  $x \in X$ . Moreover, for every  $\xi, \mu \in Y^{X \setminus \{0\}}, x \in X \setminus \{0\}$

$$\begin{aligned} & \|\mathcal{T}_m \xi(x) - \mathcal{T}_m \mu(x)\| \\ &= \left\| A\xi\left(\frac{1}{a}(m+1)x\right) + B\xi\left(-\frac{1}{b}mx\right) - A\mu\left(\frac{1}{a}(m+1)x\right) - B\mu\left(-\frac{1}{b}mx\right) \right\| \\ &\leq |A| \left\| \xi\left(\frac{1}{a}(m+1)x\right) - \mu\left(\frac{1}{a}(m+1)x\right) \right\| + |B| \left\| \xi\left(-\frac{1}{b}mx\right) - \mu\left(-\frac{1}{b}mx\right) \right\| \\ &= \sum_{i=1}^2 L_i(x) \|\xi(f_i(x)) - \mu(f_i(x))\|. \end{aligned}$$

We see that there exists  $m_0 \in \mathbb{N}$  such that

$$|A| \left| \frac{1}{a}(m+1) \right|^p + |B| \left| \frac{1}{b}m \right|^p < 1 \quad \text{for } m \geq m_0.$$

Therefore

$$\begin{aligned}
 \varepsilon^*(x) &:= \sum_{n=0}^{\infty} \Lambda_m^n \varepsilon_m(x) \\
 &= \sum_{n=0}^{\infty} c \left( \left| \frac{1}{a}(m+1) \right|^p + \left| \frac{1}{b}m \right|^p \right) \left( |A| \left| \frac{1}{a}(m+1) \right|^p + |B| \left| \frac{1}{b}m \right|^p \right)^n \|x\|^p \\
 &= \frac{c \left( \left| \frac{1}{a}(m+1) \right|^p + \left| \frac{1}{b}m \right|^p \right) \|x\|^p}{1 - |A| \left| \frac{1}{a}(m+1) \right|^p - |B| \left| \frac{1}{b}m \right|^p}, \quad x \in X \setminus \{0\}, m \geq m_0.
 \end{aligned}$$

Thus, according to Theorem 1, for each  $m \geq m_0$  there exists a unique solution  $G_m: X \setminus \{0\} \rightarrow Y$  of the equation

$$G_m(x) = AG_m \left( \frac{1}{a}(m+1)x \right) + BG_m \left( -\frac{1}{b}mx \right)$$

such that

$$\|g(x) - G_m(x)\| \leq \frac{c \left( \left| \frac{1}{a}(m+1) \right|^p + \left| \frac{1}{b}m \right|^p \right) \|x\|^p}{1 - |A| \left| \frac{1}{a}(m+1) \right|^p - |B| \left| \frac{1}{b}m \right|^p}, \quad x \in X \setminus \{0\}.$$

We show that

$$\begin{aligned}
 &\|T_m^n g(ax + by) - AT_m^n g(x) - BT_m^n g(y)\| \\
 &\leq c \left( \left| A \left| \frac{1}{a}(m+1) \right|^p + \left| B \left| \frac{1}{b}m \right|^p \right|^p \right)^n (\|x\|^p + \|y\|^p) \right) \quad (7)
 \end{aligned}$$

for every  $x, y \in X \setminus \{0\}$ ,  $n \in \mathbb{N}_0$ .

If  $n = 0$ , then (7) is simply (4). So, take  $r \in \mathbb{N}_0$  and suppose that (7) holds for  $n = r$  and  $x, y \in X \setminus \{0\}$ . Then

$$\begin{aligned}
 &\|T_m^{r+1} g(ax + by) - AT_m^{r+1} g(x) - BT_m^{r+1} g(y)\| \\
 &= \left\| AT_m^r g \left( \frac{1}{a}(m+1)(ax + by) \right) + BT_m^r g \left( -\frac{1}{b}m(ax + by) \right) \right. \\
 &\quad - A^2 T_m^r g \left( \frac{1}{a}(m+1)x \right) - AB T_m^r g \left( -\frac{1}{b}mx \right) \\
 &\quad \left. - BAT_m^r g \left( \frac{1}{a}(m+1)y \right) - B^2 T_m^r g \left( -\frac{1}{b}my \right) \right\| \\
 &\leq |A| \left\| T_m^r g \left( \frac{1}{a}(m+1)(ax + by) \right) - AT_m^r g \left( \frac{1}{a}(m+1)x \right) \right. \\
 &\quad \left. - BT_m^r g \left( \frac{1}{a}(m+1)y \right) \right\| \\
 &\quad + |B| \left\| T_m^r g \left( -\frac{1}{b}m(ax + by) \right) - AT_m^r g \left( -\frac{1}{b}mx \right) - BT_m^r g \left( -\frac{1}{b}my \right) \right\| \\
 &\leq |A| c \left( \left| A \left| \frac{1}{a}(m+1) \right|^p + \left| B \left| \frac{1}{b}m \right|^p \right|^p \right)^r \left( \left\| \frac{1}{a}(m+1)x \right\|^p + \left\| \frac{1}{a}(m+1)y \right\|^p \right) \right)
 \end{aligned}$$

$$\begin{aligned}
& + |B|c \left( |A| \left| \frac{1}{a}(m+1) \right|^p + |B| \left| \frac{1}{b}m \right|^p \right)^r \left( \left\| -\frac{1}{b}mx \right\|^p + \left\| -\frac{1}{b}my \right\|^p \right) \\
& \leq c \left( |A| \left| \frac{1}{a}(m+1) \right|^p + |B| \left| \frac{1}{b}m \right|^p \right)^{r+1} (\|x\|^p + \|y\|^p).
\end{aligned}$$

Letting  $n \rightarrow \infty$  in (7), we obtain that

$$G_m(ax + by) = AG_m(x) + BG_m(y), \quad x, y \in X \setminus \{0\}.$$

So, we have a sequence  $(G_m)_{m \geq m_0}$  of functions satisfying equation (5) such that

$$\|g(x) - G_m(x)\| \leq \frac{c \left( \left| \frac{1}{a}(m+1) \right|^p + \left| \frac{1}{b}m \right|^p \right) \|x\|^p}{1 - |A| \left| \frac{1}{a}(m+1) \right|^p - |B| \left| \frac{1}{b}m \right|^p}, \quad x \in X \setminus \{0\}.$$

It follows, with  $m \rightarrow \infty$ , that  $g$  also satisfies (5) for  $x, y \in X \setminus \{0\}$ .  $\square$

**Corollary 3.** Let  $X$  be a normed space over a field  $\mathbb{F}$ ,  $Y$  be a Banach space over  $\mathbb{K}$ ,  $a, b \in \mathbb{F} \setminus \{0\}$ ,  $A, B \in \mathbb{K}$ ,  $A + B \neq 1$ ,  $C \in Y$ ,  $c \geq 0$ ,  $p < 0$  and  $f: X \rightarrow Y$  satisfy

$$\|f(ax + by) - Af(x) - Bf(y) - C\| \leq c(\|x\|^p + \|y\|^p), \quad x, y \in X \setminus \{0\}.$$

Then  $f$  satisfies the equation

$$f(ax + by) = Af(x) + Bf(y) + C, \quad x, y \in X \setminus \{0\}.$$

*Proof.* Considering  $g(x) = f(x) + \frac{C}{A+B-1}$ ,  $x \in X$  we see that

$$\|g(ax + by) - Ag(x) - Bg(y)\| \leq c(\|x\|^p + \|y\|^p), \quad x, y \in X \setminus \{0\}.$$

So, by Theorem 2 we have

$$g(ax + by) = Ag(x) + Bg(y), \quad x, y \in X \setminus \{0\},$$

thus

$$f(ax + by) = Af(x) + Bf(y) + C, \quad x, y \in X \setminus \{0\}.$$

$\square$

**Remark 4.** If  $g$  satisfies (4) with  $A = B = 0$ , then, by (6) with  $m \rightarrow \infty$ , it follows that  $g(x) = 0$  for  $x \in X \setminus \{0\}$ .

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## References

- [1] Aoki, T.: On the stability of the linear transformation in Banach spaces. *J. Math. Soc. Jpn.* **2**, 64–66 (1950)
- [2] Badea, C.: The general linear equation in stable. *Nonlinear Funct. Anal. Appl.* **10**, 155–164 (2005)
- [3] Bourgin, D.G.: Approximately isometric and multiplicative transformations on continuous function rings. *Duke Math. J.* **16**, 385–397 (1949)
- [4] Brzdęk, J.: Hyperstability of the Cauchy equation on restricted domains. *Acta Math. Hungar.* doi:[10.1007/s10474-013-0302-3](https://doi.org/10.1007/s10474-013-0302-3)
- [5] Brzdęk, J., Chudziak, J., Páles, Zs.: A fixed point approach to stability of functional equations. *Nonlinear Anal.* **74**, 6728–6732 (2011)
- [6] Brzdęk, J., Pietrzyk, A.: A note on stability of the general linear equation. *Aequationes Math.* **75**, 267–270 (2008)
- [7] Gajda, Z.: On stability of additive mappings. *Int. J. Math. Math. Sci.* **14**, 431–434 (1991)
- [8] Găvruta, P.: A generalization of the Hyers-Ulam-Rassias stability of approximately additive mapping. *J. Math. Anal. Appl.* **184**, 431–436 (1994)
- [9] Hyers, D.H.: On the stability of the linear functional equation. *Proc. Natl. Acad. Sci. USA* **27**, 222–224 (1941)
- [10] Hyers, D.H., Isac, G., Rassias, Th.M.: *Stability of Functional Equations in Several Variables*. Birkhäuser, Berlin (1998)
- [11] Jung, S.M.: *Hyers-Ulam stability of functional equations in mathematical analysis*. Hadronic Press, Palm Harbor (2001)
- [12] Kuczma, M.: *An introduction to the theory of functional equation and inequalities*. PWN, Warszawa (1985)
- [13] Maksa, Gy., Páles, Zs.: Hyperstability of a class of linear functional equations. *Acta Math. Acad. Paedag. Nyiregyháziensis* **17**, 107–112 (2001)
- [14] Popa, D.: Hyers-Ulam-Rassias stability of the general linear equation. *Nonlinear Funct. Anal. Appl.* **7**, 581–588 (2002)
- [15] Rassias, Th.M.: On the stability of the linear mapping in Banach spaces. *Proc. Am. Math. Soc.* **72**, 297–300 (1978)
- [16] Rassias, Th.M.: On a modified Hyers-Ulam sequence. *J. Math. Anal. Appl.* **158**, 106–113 (1991)

Magdalena PiszczeK  
Institute of Mathematics  
Pedagogical University  
Podchorążych 2  
30-084 Kraków  
Poland  
e-mail: magdap@up.krakow.pl

Received: March 1, 2013

Revised: April 18, 2013